

CONTENTS

ORIGINAL STUDIES

<i>Is the level of vitamin D deficiency correlated with the severity and bilaterality in slipped capital femoral epiphysis? A case series study</i> , Shady Elbeshry et al.	217
<i>Long-term outcome of extraarticular subtalar arthrodesis in children with cerebral palsy using modified Grice technique</i> , Jiri Jochymek, Tereza Peterkova	223
<i>Assessment of early Achilles tenotomy in the newborn idiopathic pes equinovarus</i> , Ersin Mutlu et al.	231
<i>Reliability of radiological measurements and treatment recommendations of displaced paediatric humeral medial epicondyle fractures</i> , Christoph Arneitz et al.	237
<i>Pediatric distal humeral supracondylar fracture – achievement of optimal pinning configuration</i> , Han Hong Chong, Assad Qureshi	245
<i>Quantitative gait analysis in children with osteogenesis imperfecta: relationship between gait deviations and clinical features</i> , Solange De Wouters et al.	255
<i>Open reduction in treatment of neglected elbow dislocation in children</i> , Ahmed El Mouloua et al..	263
<i>The management of orthopedic trauma surgery during the COVID-19 pandemic in Turkey</i> , Ortac Gurcan et al.	269
<i>The effect of transitional vertebrae and spina bifida occulta on disc herniation, disc degeneration, and end-plate changes in pediatric patients with low back pain</i> , Ozge Gulsum Illeez et al.	275
<i>Zero-profile implant versus integrated cage-plate implant in treatment of single level cervical disc disease</i> , Mohammed Zayan Ibrahim et al.	285
<i>Impact of job characteristics on return-to-work interval following arthroscopic partial meniscectomy</i> , Jef Van Doninck et al.	293
<i>The survivorship of revision total hip replacement with severe proximal bone deficiency using 15 modular taper fluted prosthesis</i> , Siddharth Govilkar et al.	303
<i>Hip fracture specialists facilitate low-dose spinal anaesthesia in fractured neck of femur surgery</i> , David E. Brooks et al.	311
<i>Total knee arthroplasty for acute tibial plateau fractures: a survey amongst 68 Dutch orthopaedic surgeons</i> , Mees K. Hesmerg et al.	319
<i>Mid-term outcomes of posterior capsular release for fixed flexed deformity after total knee arthroplasty</i> , Theofylaktos Kyriakidis et al.	329
<i>The addition of oral Tranexamic acid to knee arthroplasty patients does not further improve blood loss: a double blinded randomized control trial</i> , Ciaran McDonald et al.	335
<i>Tibial cancellous bone auto-grafting for medial open-wedge high tibial osteotomy: bone void filling with tissue harvested from osteotomized medullary canal</i> , Cumhur Cevdet Kesemenli et al.	342
<i>Quadriceps tendon vs hamstring autograft in primary ACL reconstruction – a comparative study with minimum two-year follow-up</i> , Maria Victoria Pomenta Bastidas et al.	347
<i>Are running stitches needed in DT4 Single Anteromedial Bundle Biological Augmentation transplant preparation ?</i> , Maxime Huard et al.	355
<i>Comparison of the WALANT and the PFNB techniques in the surgical treatment of unimalleolar fractures: a prospective study</i> , Ceyhun Çağlar et al.	359

<i>Non-contact wound area assessment by digital planimetry using photo editing software</i> , Pradyumna Krishna Majumdar <i>et al.</i>	368
<i>Instability after reverse total shoulder arthroplasty: risk factors and how to avoid them</i> , Lorena Pena <i>et al.</i>	372
<i>Radial head fractures: a quantitative analysis</i> , Silvio Lampaert <i>et al.</i>	380
<i>The prevalence of frozen shoulder in patients with limited Dupuytren disease</i> , Maarten Van Nuffel <i>et al.</i>	387
<i>Radial tuberosity anatomy in intramedullar repair of distal biceps tendon ruptures. A radiological study</i> , Laurens Van Melkebeke <i>et al.</i>	392
<i>Treatment practice for Dupuytren disease in Belgium before 2020: results from an online survey</i> , Maarten Van Nuffel <i>et al.</i>	399
<i>Outcome of the MatOrtho arthroplasty for PIP osteoarthritis with a minimum follow-up of two years</i> , Kathleen Janssens, Frederik Verstreken	410