



External to internal glenohumeral strength ratio in non-traumatic rotator cuff pathologies

Peter VERSPEELT¹, Martine DE MUYNCK², Guy VANDERSTRAETEN², Luc VANDEN BOSSCHE², Gaetane STASSIJNS³, Levent ÖCZAKAR⁴

¹Department of Physical Medicine and Rehabilitation, Antwerp University Hospital and Ghent University Hospital, Antwerp and Ghent, Belgium; ²Faculty of Medicine and Health Sciences, Ghent University, Department of Physical Medicine and Rehabilitation, University Hospital Ghent, Ghent, Belgium; ³Faculty of Medicine and Health Sciences, University of Antwerp, Department of Physical Medicine and Rehabilitation, Antwerp University Hospital, Antwerp, Belgium; ⁴Department of Physical and Rehabilitation Medicine, Hacettepe University Medical School, Ankara, Turkey

Correspondence at: Verspeelt Peter, Department of Physical Medicine and Rehabilitation, University Hospital Antwerp, Berchemboslaan 13, 2600 Berchem, Belgium, E-mail: peter.verspeelt@gmail.com

Background and study aims: A correct agonist -antagonist strength relationship for shoulder external and internal rotation is necessary for functional stability of the shoulder. This strength relationship is described by the ratio of external to internal strength (ER/IR). The aim of this study is to produce comparative data as regards the ER/IR ratio in subjects with different non-traumatic rotator cuff diseases. **Design and setting:** A cross-sectional study in an outpatient clinic in a tertiary care university hospital. **Methods:** In 55 subjects with rotator cuff disease (confirmed by physical examination and assessed by ultrasound and magnetic resonance arthrography), the ER/IR ratio of the shoulder was isometrically measured with a hand-held dynamometer and compared with values pertaining to the unaffected shoulder of the same individuals. **Results:** The mean ER/IR values in the overall group were 0.89 (SD 0.18) and 0.94 (SD 0.22) for the affected and unaffected shoulders, respectively. The ratio was 0.87 (SD 0.23) in patients with subdeltoid bursitis, 0.88 (SD 0.16) in rotator cuff tendinopathy and 0.87 (SD 0.22) in patients with rotator cuff tears. **Conclusions:** The ER/IR ratio appears to be similar between the affected and unaffected shoulders of subjects with nontraumatic cuff pathologies.

Key words: Shoulder, rotator cuff, isometric strength, dynamometry, ultrasound.

INTRODUCTION

The glenohumeral joint maintains the largest range of motion among all the body joints and this requires efficiency of every shoulder muscle. An appropriate balance of agonist and antagonist muscle strength is necessary to provide sufficient active stability to maintain normal shoulder kinematics. Likewise, any disorder in the balance of these muscles will negatively influence the shoulder biomechanics and increase the possibility of shoulder disease.

Rotator cuff (RC) disorders are the underlying cause in 65% to 70% of people with shoulder pain^{1,2}. Several studies have demonstrated that an imbalance between the strength of the external rotator (ER) and internal rotator (IR) muscles is present in subjects with RC disorders^{3,4,5,6,7,8,9,10}. However, in many of those studies either the sample size was small or lacks data from the contralateral side. Besides many of these studies lack the necessary imaging such as ultrasound and MRI puts limitations to the results of these studies. This study

tried to avoid these pitfalls using a greater sample size, different gender and age groups and strong additional imaging such as ultrasound and MRI of the affected shoulder and ultrasound of the control shoulder.

Accordingly, the aim of this study was to comparatively measure the ER/IR values in patients with various RC disorders.

MATERIALS AND METHODS

Subjects with shoulder pain for more than 12 weeks were recruited if physical examination findings such as a painful arc, a positive Hawkins-Kennedy and Jobe test were encountered. Patients were excluded if they had previous shoulder trauma, frozen shoulder, inflammatory arthritis and radiographic evidence of osteoarthritis. All included subjects underwent bilateral shoulder ultrasound (US) examination and unilateral magnetic resonance arthrography (MRA) for the symptomatic shoulders.

Depending of the US and MRA findings, patients were divided into three groups according to the presence of bursitis, RC tendinopathy or RC partial/full-thickness tears. Patients were excluded when two or more findings were simultaneously present. As the US examination was performed on both shoulders, data from subjects who had abnormal findings in their asymptomatic shoulders were not used in the study. 55 patients with nontraumatic unilateral RC disease were included in the study. This study was approved by the institutional medical ethics review board. All participants signed an informed consent form before participation.

The US was performed using a Toshiba Aplio 300 using a 7-9 MHz transducer following the EURO-MUSCULUS/USPRM protocols¹¹. MRA of the symptomatic shoulder was performed on a Ingenia Elition 3.0T with a large shoulder coil. The US and MRA were performed in the same university radiology service on the same day and evaluated by two MD's specialized in musculoskeletal radiology and holding a PhD degree and each with more than 15 years of experience.

Isokinetic and isometric testing are valid methods for measurement of muscle strength providing an unbiased estimation of muscle strength using a linear scale enabling accuracy and sensitivity. Isometric strength testing with a hand held dynamometer (HHD) is a relatively inexpensive and efficient way to assess strength^{12,13,14}. In many studies HHD has proven to have good-to-excellent intra- and interrater reliability and also good validity^{15,16,17,18,19,20,21,22,23,24}. A HHD (Mecmesin myometer, Broadridge Heath, England) with a measurement range of 0 to 300 Newton (N) and a precision of 0.1 was used. The HHD was factory calibrated and recalibrated before each use. In measuring isometric strength, the "make" method rather than the "break" method was used given the higher reliability coefficients^{22,26}. After the examiner explained the test and showed the direction of movement, subjects were asked to build up their force gradually to a maximum voluntary effort over a two second period and to apply maximum force for about three seconds. Each task was repeated twice with 15 seconds between measurements. The peak isometric strength was the average of the two trials and was used for analysis.

During the testing, subjects were in standing position with their arms besides the body. The testing order was standardized and consisted of abduction of the shoulder, flexion of the elbow and external and internal rotation of the shoulder. For testing the abduction, the shoulder was at neutral and the elbow in extended position. The dynamometer was placed proximal to the styloid process. For elbow flexion, the shoulder was at neutral, the elbow at 90° flexion and the forearm in supination. The dynamometer was placed proximal to the styloid process. For internal rotation, shoulder was at neutral and the elbow at 90° flexed position. The dynamometer was placed just proximal to the styloid process. For external rotation, the shoulder was at 0° abduction and 45° internal rotation²⁷.

To minimize bias, all tests were performed by the same rehabilitation physician with long experience in muscle testing. The strength of the examiner was strong enough to hold against the isometric contraction of the subjects being tested.

All statistical analysis was performed using SPSS Statistics 22 Data are presented as median (range) and mean (standard deviation) values. Normality of the data was tested by Shapiro-Wilk test and visual inspection of the QQ-plots. As most of the ER/IR ratios were not normally distributed, non-parametric tests were used. ER/IR ratios were compared using Wilcoxon signed rank and Kruskal-Wallis tests for affected vs. unaffected shoulders and for shoulders among three groups, respectively. A p-value of 0.05 or lower was considered significant.

RESULTS

Demographic data of the participants are summarized in Table I. Depending of the US and MRA findings, the patients were categorized into three groups (based on causality) as bursitis (n=10), cuff tendinopathy (n=27) and cuff tears (n=18). ER/IR ratios of the individuals are given in Table II. The ER/IR ratios of the affected (0.89±0.18, range 0.43-1.24) and unaffected (0.94±0.22, range 0.22-1.86) shoulders were similar (p=0.11). The subjects with bursitis showed an ER/IR ratio of 0.87±0.23 in the affected and 0.98±0.18 in the unaffected shoulder. The subjects with tendinopathy showed an ER/IR ratio of 0.88±0.16 in the affected and,

Table I. — Demographic data of the subjects based on gender and age category.

	n	%	Average age	20-40y	40-60y	60+y
Male	37	(67.3%)	47.2 (21-69)	n=11 (20.0%)	n=17 (30.9%)	n=9 (16.4%)
Female	18	(32.7%)	50.6 (26-65)	n=3 (5.5%)	n=13 (23.6%)	n=2 (3.6%)

Table II. — ER/IR ratio in 55 subjects.

		Ratio ER/IR				Ratio ER/IR	
Subject	Gender	Sympt	Asympt	Subject	Gender	Sympt	Asympt
1	M	0.86	0.99	29	M	0.94	0.88
2	F	0.89	1.13	30	F	0.83	0.90
3	F	1.14	1.04	31	M	1.22	1.03
4	M	0.52	1.11	32	F	1.14	0.88
5	M	0.87	1.17	33	M	1.03	0.82
6	M	0.89	0.89	34	M	0.97	1.34
7	M	1.21	1.00	35	F	0.92	0.79
8	F	0.91	0.87	36	M	1.02	0.77
9	F	0.43	1.39	37	M	1.07	1.19
10	F	0.55	0.91	38	M	0.64	0.83
11	M	1.08	0.93	39	F	0.74	0.63
12	M	0.91	0.75	40	M	0.82	0.93
13	M	0.84	1.20	41	M	0.61	0.22
14	F	0.88	0.77	42	M	0.81	0.90
15	F	0.92	1.08	43	M	1.23	0.90
16	M	0.90	0.94	44	M	0.76	1.02
17	F	0.81	0.96	45	M	0.70	1.01
18	F	0.81	1.03	46	M	1.09	1.86
19	F	1.00	0.58	47	F	0.75	0.78
20	M	1.24	0.98	48	M	0.75	0.85
21	M	0.90	1.00	49	F	0.76	0.82
22	M	0.98	0.97	50	M	0.76	0.99
23	M	1.04	0.95	51	M	0.73	0.87
24	M	0.94	1.02	52	M	0.94	1.10
25	M	0.95	0.75	53	F	0.79	0.80
26	M	1.08	1.03	54	M	0.79	0.73
27	M	0.84	0.82	55	M	0.66	0.66
28	F	0.85	1.01				

Table III. — Average, standard deviation and p-value for Bursitis, Cuff tendinosis and Cuff Tear.

	Sympto		Asympto		p
	average	stand. Dev.	average	stand. Dev.	
Bursitis	0,87	0,23	0,98	0,18	0,33
Cuff tendinosis	0,88	0,16	0,93	0,27	0,29
Cuff tear	0,93	0,16	0,86	0,08	0,17

0.93±0.27 in the unaffected shoulder. The subjects with a cuff tear showed an ER/IR ratio of 0.87±0.22 in the affected and, 0.95±0.19 in the unaffected shoulder (see

Table III). The difference between the affected vs. the unaffected shoulder in the 3 causalities was statistically insignificant, with all p > 0.05

DISCUSSION

A lot of research has already been done on measuring muscle strength of the shoulder in overhead athletes and healthy people. Less research has been done on the strength in cohorts with non-traumatic RC disease. We systematically searched electronic databases with the following search term: muscle strength of the glenohumeral muscles in a non athletic population with non-traumatic RC disease. Reports on shoulders with instability or SLAP lesions were omitted. As in the case with any literature review some evidence may have been unintentionally overlooked. Many studies have a full data set of strength measurements but do not discuss the ER/IR ratio^{28,29,30,31,32,33,34,35,36}. Only one study comments on the ER/IR ratio whereby 51 cases with subacromial shoulder impingement and 51 asymptomatic controls matched for age, gender and hand dominance were evaluated using isokinetic testing³⁷. The authors report that either using eccentric or concentric testing the ER/IR were similar between the affected and nonaffected shoulder. We observed that the ER/IR ratios are similar between affected and nonaffected shoulders so our results confirm the findings of the above mentioned study. In the overall group the ER/IR ratio is 0.89 (SD 0.18) for the affected shoulder and 0.94 (SD 0.22) for the unaffected shoulder. This difference is not significant as $p > 0.05$.

When sub analyzing the group for the 3 different pathologies the results are: In the bursitis group for the affected shoulder 0.87 (SD 0.23) and the nonaffected shoulder 0.98 (SD 0.18), in the tendinopathy group for the affected shoulder 0.88 (SD 0.16) and the nonaffected shoulder 0.93 (SD 0.27) and in the cuff tear group the affected shoulder 0.93 (SD 0.16) and the nonaffected shoulder 0.86 (SD 0.08). These differences are non-significant with $p = 0.33$ in the bursitis group, $p = 0.29$ in the cuff tendinopathy group and $p = 0.17$ in the cuff tear group.

The strengths of the study are the following. First we measured the ER/IR ratio in the affected and the unaffected shoulder in the same person. This leads to results with more sustainability. Second by examining each affected shoulder with ultrasound and MRA we were completely sure about the underlying pathology and excluding any posttraumatic pathology. Third by examining all nonaffected shoulders with ultrasound we excluded any underlying pathology in this shoulder.

There are some limitations worth noting for the study. First, the data generated by muscle testing must be cautiously handled – also taking in account the possible impact of discomfort or pain during the testing.

We tried to accommodate the subjects in the most appropriate position in this regard. Second, although upper limb dominance is negligible in normal subjects in contrast to overhead athletes, the lack of pertinent analysis might be another limitation in the study^{38,39}. Third the sample size of our study was sometimes small with regards to subgroup analysis.

CONCLUSION

This study concludes that the ER/IR ratio is not different in the with a RC disease affected shoulder and the contralateral not affected shoulder. In planning the rehabilitation of shoulder dysfunction, this ratio seems to be of no great importance. There is need for future studies on assessing other shoulder strength ratios such as abduction/external(ABD/ER) or abduction/ internal (ABD/IR).

Conflict of interest: the authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

The authors report no involvement in the research by any sponsor that could have influenced the outcome of this work.

Authors' contributions: Author P Verspeelt has given substantial contributions to the conception and the design of the manuscript and to acquisition, analysis and interpretation of the data. All authors have participated to drafting the manuscript, author L Oczakar revised it critically. All authors read and approved the final version of the manuscript.

REFERENCES

1. Chard MD, Hazleman R, Hazleman BL, King RH, Reiss BB. Shoulder disorders in the elderly: a community survey. *Arthritis Rheum.* 1991;34 (6):766-769.
2. Vecchio P, Kavanagh R, Hazleman BI, King RH. Shoulder pain in a community-based rheumatology clinic. *Br J Rheumatol.* 1995;34 (5):440-442.
3. Byram IR, Bushnell BD, Dugger K et al. Preseason shoulder strength measurements in professional baseball pitchers: identifying players at risk for injury. *Am J Sports Med.* 2010; 38:1375-1382. <https://doi.org/10.1177/0363546509360404>.
4. Clarsen B, Bahr R, Andersson SH et al. Reduced glenohumeral rotation, external rotation weakness and scapular dyskinesis are risk factors for shoulder injuries among elite male handball players: a prospective cohort study. *Br J Sports Med.* 2014;48:1327-1333. <https://doi.org/10.1136/bjsports-2014-093702>.
5. Dauty M, Delbrouck C, Huguet D, et al. Reproducibility of concentric and eccentric isokinetic strength of the shoulder rotators in normal subjects 40 to 55 years old. *Isokinet Exerc Sci.* 2003;(11):95-100. <https://doi.org/10.3233/IES-2003-0134>.
6. Edouard P, Frize N, Calmels P, et al. Influence of rugby practice on shoulder internal and external rotators strength. *Int J Sports Med.* 2009;(30):863-867. <https://doi.org/10.1055/s-0029-1237391>.

7. Ellenbecker TS, Davies GJ. The application of isokinetics in testing and rehabilitation of the shoulder complex. *J Athl Train*. 2000;(35):338-350.
8. Leroux JL, Codine P, Thomas E, Pocholle M, Mailhe D, Blotman F. Isokinetic evaluation of rotational strength in normal shoulders and shoulders with impingement syndrome. *Clin Orthop Relat Res*. 1994;jul (304):108-15.
9. Leroux T, Chahal J, Wasserstein D, Verma NN, Romeo AA. A systematic review and meta-analysis comparing clinical outcomes after concurrent rotator cuff repair and long head biceps tenodesis or tenotomy. *Sports Health*. 2015;7:303-307. <https://doi.org/10.1177/1941738114539627>.
10. Warner JJ, Micheli LJ, Arslanian LE, et al. Patterns of flexibility, laxity, and strength in normal shoulders and shoulders with instability and impingement. *Am J Sports Med*. 1990;(18):366-375.
11. Ozcakar L, Kara M, Wang TG, De Muynck M. Euromusculus/USPRM Basic scanning protocols: a practical guide for physiatrists. *Eur J Phys Rehabil Med*. 2015;51:477-478. PMID: 26173449.
12. Byl NN, Richards S, Asturias J. Intrarater and interrater reliability of strength measurements of the biceps and deltoid using a hand held dynamometer. *J Orthop Sports Phys Ther*. 1988;9 (12):395-398.
13. Rabin SI, Post Ma. Comparative study of clinical muscle testing and cybex evaluation after shoulder operations. *Clinical Orthopedics*. 1990;258:147-156.
14. Sullivan JS, Chesley A, Herbert G, Mc Faull S, Scullion D. The validity and reliability of hand -held dynamometer in assessing isometric external rotator performance . *J Orthop Sports Phys Ther*. 1988;10:213-217.
15. Baschung Pfister P, de Bruin ED, Sterkele I, Maurer B, de Bie RA, Knols RH. Manual muscle testing and hand-held dynamometry in people with inflammatory myopathy: an intra- and interrater reliability and validity study. *PLoS ONE*. 2018;29 (13(3)). <https://doi.org/10.1371/journal.pone.0194531>.
16. Cools AM, De Wilde L, Van Tongel A, Ceysens C, Ryckewaert R, Cambier DC. Measuring shoulder external and internal rotation strength and range of motion: comprehensive intrarater reliability study of several testing protocols. *J Shoulder Elbow Surg*. 2014;23:1454-1461. <https://doi.org/10.1016/j.jse.2014.01.006>.
17. Fieseler G, Laudner KG, Irlenbusch L, et al. Inter- and intrarater reliability of goniometry and hand held dynamometry for patients with subacromial impingement syndrome. *Journal of Exercise Rehabilitation*. 2017;13 (6):704-710 <https://doi.org/10.12965/jer.1735110.0555>.
18. Harbo T, Brincks J, Andersen H. Maximal isokinetic and isometric muscle strength of major muscle groups related to age, body mass, height and sex in 178 healthy subjects. *Eur J Appl Physiol*. 2012;112 (1):267-275. <https://doi.org/10.1007/s.00421-011-1975-3>.
19. Hayes K, Walton JR, Szomor ZL, Murrell GA. Reliability of 3 methods for assessing shoulder strength. *J Shoulder Elbow Surg*. 2002;11 (1):33-39 <https://doi.org/10.1067/mse.2002.119852>.
20. Kim HM, Teefey SA, Zelig A, Galatz LM, Keener JD, Yamaguchi K. Shoulder strength in asymptomatic individuals with intact compared with torn rotator cuffs. *J Bone Joint Surgery Am*. 2009;91 (2):289-296. <https://doi.org/10.2106/jbjs.h.00219>.
21. Kolber MJ, Beekhuizen K, Ming-Shun S. Cheng, Fiebert IM. The reliability of hand-held dynamometry in measuring isometric strength of the shoulder internal and external rotator musculature using a stabilization device. *Physiotherapy Theory and Practice*;2007;23 (iss 2):161-8. <https://doi.org/10.1080/09593980701213032>.
22. Magnusson SP, Gleim GW, Nicholas JA. Subject variability of shoulder abduction strength testing. *Am J Sports Med*. 1990;18:349-353.
23. Rieman BL, Davies GJ, Ludwig L, Gardenhour H. Hand-held dynamometer testing of the internal and external rotator musculature based on selected positions to establish normative data and unilateral ratios. *J Shoulder Elbow Surg* . 2010;19 (8):1175-1183. <https://doi.org/10.1016/j.jse.2010.05.021>.
24. Roy JS, MacDermid JC, Orton B, et al. The concurrent validity of a hand -held versus a stationary dynamometer in testing isometric shoulder strength. *J Hand Ther*. 2009;oct-dec 22 (4):320-6. <https://doi.org/10.1016/j.jht.2009.04.008>.
26. Stratford PW, Balsor BE. A comparison of make and break tests using a hand-held dynamometer and the Kin-Com. *J Orthop Sports Phys Ther*. 1994;19 (1):28-32.
27. Kelly BT, Kadrmas, WR, Speer KP. The manual muscle examination for muscle strength: an electromyographic investigation. *Am J Sports Med*. 1996;24 (5):581-588.
28. Akyol Y, Ulus Y, Durmus D, et al. Shoulder muscle strength in patients with subacromial impingement syndrome: its relationship with duration of quality of life and emotional status. *Turk J Phys Med Rehab*. 2013;59:176-181. <https://doi.org/10.4274/tfll.59837>.
29. Bak K, Magnusson SP. Shoulder strength and range of motion in symptomatic and pain free elite swimmers. *Am J Sports Med*. 1997;25 (4):454-9.
30. Letafatkar A, Abbaszadeh Ghanati H, Sheikhi B. Patients with subacromial impingement syndrome exhibit altered shoulder rotor muscles eccentric and concentric peak torque. *Journal of Rehabilitation Sciences and Research*. 2018;5:99-105.
31. Malerba JL, Adam ML, Harris BA, Krebs DE . Reliability of dynamic and isometric testing of shoulder external and internal rotators. *J Orthop Sports Phys Ther*. 1993;18 (4):543-52. <https://doi.org/10.2519/jospt.1993.18.4.543>.
32. McClure PW, Michener LA, Karduna AR. Shoulder function and 3- dimensional scapular kinematics in people with and without impingement syndrome. *Physical Therapy*. 2006;86 (8):1075-1090. <https://doi.org/10.1093/ptj/86.8.1075>.
33. McLaine SJ, Ginn KA, Fell JW, Bird ML. Isometric shoulder strength in young swimmers. *Journal of Science and Medicine in Sport*. 2018;21:35-39. <https://doi.org/10.1016/j.jsams.2017.05.003>.
34. Saccol MF, Zanca GG, Ejnisman B, de Mello MT, Mattiello SM. Shoulder rotator strength and torque steadiness in athletes with anterior shoulder instability or SLAP lesion. *J Science Med Sport*. 2014;17 (5):463-8. <https://doi.org/10.1016/j.jsams.2013.10.246>.
35. Stickley CD, Hetzler RK, Freemyer BG, Kimura IF. Isokinetic peak torque ratios and shoulder injury history in adolescent female volleyball athletes. *Journal of Athletic Training*. 2008;43 (6):571-577. <https://doi.org/10.4085/1062-6050-43.6.571>.
36. Wilbanks SR, C. S Bickel . Scapular stabilisation and muscle strength in manual wheelchair users with spinal cord injury and subacromial impingement. *Top Spinal Cord Inj Rehabil*. 2016;22 (1):60-70. <https://doi.org/10.1310/sci2201-60>.
37. Land H, Gordon S, Watt K. Isokinetic clinical assessment of rotator cuff strength in subacromial shoulder impingement. *Musculoskeletal Science and Practice*. 2017;27:32-39. <https://doi.org/10.1016/j.msksp.2016.11.012>.
38. Ellenbecker TS, Mattalino AJ. Concentric isokinetic shoulder internal and external rotation strength in professional baseball pitchers. *J Orthop Sports Phys Ther*. 1997;(25):323-328.
39. Hinton RY. Isokinetic evaluation of shoulder rotational strength in high school baseball pitchers. *Am J Sports Med*. 1988;(16):274-279.